

Fe K- α Emission from a Decaying Magnetar Model of Gamma-Ray Bursts

M.J. Rees ¹ & P. Mészáros ^{1,2,3}

¹Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, U.K.

²Dpt. of Astronomy & Astrophysics, Pennsylvania State University, University Park, PA 16803

³Institute for Theoretical Physics, University of California, Santa Barbara, CA 93106-4030

ABSTRACT

The recent report of X-ray Fe features in the afterglow of the gamma-ray burst GRB 991216 may provide important clues for identifying the nature of its progenitor and constraining the burst mechanism. We argue that the strong line emission can be attributed to the interaction of a continuing (but decaying) post-burst relativistic outflow from the central engine with the progenitor stellar envelope at distances less than a light-hour. Only a small mass of Fe is then required, which could have been readily produced by the star itself.

Subject headings: Gamma-rays: Bursts - X-rays - Cosmology: Miscellaneous

1. Introduction

Recently, Fe K- α and K-edge X-ray features have been reported with high significance in the afterglow of the bright burst GRB 991216 and GRB 000214 after about 1 day (Piro, *et al.*, 2000, Antonelli et al, 2000). A straightforward interpretation of this observation would imply a mass $\gtrsim 10^{-2} - 1M_{\odot}$ of Fe at a distance of about one light-day, possibly due to a remnant of an explosive event or supernova which occurred days or weeks prior to the gamma-ray burst itself.

Here, we suggest an alternative and perhaps less restrictive scenario: an extended, possibly magnetically dominated wind from a GRB impacting the expanding envelope of a massive progenitor star. This could be due either to a spinning-down millisecond super-pulsar or to a highly-magnetised torus around a black hole (e.g. Wheeler, *et al.* 2000), which could produce a luminosity that was still, one day after the original explosion, as high as $L \sim 10^{47} \text{ erg s}^{-1}$. This luminosity may not dominate the continuum afterglow; but we argue that it could be efficiently reprocessed, by a modest amount $M_{Fe} \sim 10^{-8} M_{\odot}$ of material at distances $\lesssim 10^{13} \text{ cm}$, into an Fe line luminosity comparable to the observed value, together with a contribution to the X-ray continuum. Under this interpretation, the dominant continuum flux in the afterglow, even in the X-ray band, is still attributable to a standard decelerating blast wave.

2. Energy Input and Fe Line Strength

The typical GRB model assumes that the energy input episode is brief, typically $t_b \lesssim 1 - 10^2$ s, its energy and mass outflow being either a delta or a top-hat function. However, peculiarities in the early stages of some afterglows, e.g. GRB 970508, have served as motivation for considering a more extended input period in which the energy or mass outflow rate may vary in time and the late energy input could exceed the prompt contribution (e.g. Rees & Mészáros 1998, Sari & Mészáros 2000). Here we consider a related model, in which the power output continues at a diminishing rate, for a longer time of hours to days. The prolonged activity could arise if orbiting debris around a newly-formed black hole takes a long time to be completely swallowed, or if the central object becomes a fast-spinning neutron star, rather than a black hole. Collapsar or hypernova models (e.g. MacFadyen & Woosley, 1999; Paczyński, 1998; Woosley 1993) or magnetar-like GRB models (Usov, 1994, Thompson, 1994, Wheeler *et al.* 2000) provide a natural scenario for a sudden burst followed by a more slowly decaying energy input.

The power output would be primarily in a magnetically-driven relativistic wind (which, even during the later phases discussed here, would be hugely super-Eddington). The generic magnetized outflow from a spinning compact object is

$$L_m \sim 2\pi r_o^2 c (B^2/4\pi) \sim 1.5 \times 10^{52} r_{o6}^2 B_{15}^2 \text{erg s}^{-1} \sim 1.5 \times 10^{47} r_{o6}^2 B_{12.5}^2 \text{erg s}^{-1}, \quad (1)$$

where the suffix denotes the B-field in gauss. The characteristic radius $r_o = 10^6 r_{o6}$ cm could be the inner radius of the accretion torus in a BH model, or, in a superpulsar model spinning with near-breakup angular velocity, the radius of the light cylinder. In a normal pulsar model, the field is assumed to maintain a steady value, and the luminosity declines as the spin rate slows down. However, during the early stages, B might decline more rapidly than the slowing-down timescale: the power output then declines in proportion to B^2 . For instance, if a torus is losing angular momentum due to magnetic torques, and is gradually draining into the hole, the magnetic field (and therefore the associated MHD torques and wind-driven energy losses) would gradually decline in step with the surviving mass. Alternatively, if the central object were a rapidly-spinning neutron star, the field may decline as uniform rotation is established and the higher moments decay away.

There is no reason to expect that the field decay would follow a power law. However, just as an illustration, we note, from equation (1), that a luminosity decay $L(t) \propto t^{-1.25}$, which would lead to 10^{47} erg/s after one day, could be a consequence of a drop in B from 10^{15} G to 3×10^{12} G in a compact structure with stored energy of at least 10^{52} ergs whose characteristic spin period remained constant (at a fraction of a millisecond).

A similar argument could be developed based on the concept of α -viscosity. For a hot dense torus around the BH resulting from collapse of the core of the progenitor star, the α -viscosity parameter is $\alpha \sim B^2/(4\pi\rho v_s^2) \sim 0.05 B_{15}^2 \rho_{13}^{-1} T_9^{-1}$, and the viscous accretion time for a torus of outer radius $10^9 r_9$ cm is $t_{visc} \sim (\alpha\Omega)^{-1} \sim 10 r_9^{3/2} B_{15.5}^{-2} \rho_{13} T_9$ s, while for a lower field of 3×10^{12} G

it is $t_{visc} \sim 1.5 r_9^{3/2} B_{12.5}^{-2}$ days for $nT \sim \text{constant}$, and the accretion of $\lesssim 10^{-2} M_\odot$ in $t \sim 1$ day is sufficient to provide a characteristic $L \sim 10^{47} \text{ erg s}^{-1}$ at a day.

We envisage that the burst is triggered by the collapse of a core within a massive stellar envelope, as in the scenario developed by Woosley and his collaborators. A funnel along the rotation axis would have been blasted open during the 1-100 s duration of the original burst; it would subsequently enlarge, owing to the post-explosion expansion of the envelope of the progenitor star (e.g. Eichler & Levinson 1999; Woosley 1993). The ram pressure of the continuing MHD outflow would further enlarge the funnel and could, after one day, have expelled the envelope material from a region 10^{13} cm across, even in the equatorial plane (this would require velocities of no more than 10^8 cm/s).

The magnetised wind from the compact remnant (which we assume to be relativistic) would develop a stand-off shock before encountering the envelope material, and shocked relativistic plasma would be deflected along the funnel walls. In the absence of magnetic fields, the contact discontinuity between the shocked jet and stellar gas would have a tendency to develop Kelvin-Helmholtz instabilities, which could lead to bulk heating of the stellar material. This would extend to large Thomson depths $\tau_T \gg 1$, required to ensure a fluid behavior leading to non-magnetic K-H instabilities. However, magnetic fields are expected, since even if initially absent the instabilities would lead to mixing with the highly magnetized jet material. The K-H equivalent of highly relativistic MHD oblique shear instabilities is poorly understood, and it is unclear how deeply, if at all, such bulk heating would penetrate the thermal material outside the funnel wall. However, for energy deposition spread over a layer thick enough to have $\tau_T \gg 1$, the cooling rate (due to comptonisation, bremsstrahlung and recombination) would be high enough to reduce the temperature of the bulk-heated electrons and protons to the equivalent black body temperature, estimated as $T_e \sim T_p \sim (L_m / \sigma 4\pi r_*^2)^{1/4} \sim \text{few} 10^6 L_{47} r_{13}^{-1/2}$ K.

A more efficient heating mechanism of the stellar funnel wall gas is radiative heating, which would deposit energy within shallower layers with modest scattering optical depth. Non-thermal electrons are expected to be accelerated behind the standoff shock in the jet material; the transverse magnetic field strength (which decreases as $1/r$ in an outflowing wind) would be of order 10^4 G at 10^{13} cm – strong enough to ensure that the shock-accelerated electrons cool promptly, yielding a power-law continuum extending into the X-ray band. Some of these X-rays would escape along the funnel, but at least half (the exact proportion depending on the geometry and flow pattern) would irradiate the material in the stellar envelope. For the high radiative efficiencies expected in relativistic shocks, the inward-directed radiative heat flux would be comparable to that of a bulk heat flux from instabilities. However since the radiative flux is deposited in shallower layers containing less mass than the bulk heat flux, the radiatively heated shallow layer would be expected to be substantially hotter than a deeper bulk-heated region, its temperature being determined by photoionization equilibrium.

The shocked relativistic material in the jet MHD flow exerts a pressure that is comparable

with the ram pressure of the pre-shocked outflow. This is $\alpha L_m/4\pi r^2 c$, where $r = 10^{13} r_{13}$ is the distance of the funnel wall, and $\alpha \lesssim 1$ is a geometrical factor. This must be balanced by the thermal pressure $3n_e kT$ in the photoionized layers of the stellar envelope forming the outside the wall of the funnel, giving a characteristic electron density

$$n_e = \alpha L_m/6\pi r^2 c kT \sim 10^{17} \alpha L_{47} r_{13}^{-2} T_8^{-1} \text{ cm}^{-3} \quad (2)$$

The temperature parametrization used here is consistent with the range expected from photoionization equilibrium. The recombination time for hydrogenic Fe in the funnel walls photoionized by the non-thermal continuum produced by the stand-off shock is

$$t_{rec} = 6 \times 10^{-6} T_8^{1/2} n_{17}^{-1} = 6 \times 10^{-6} \alpha^{-1} L_{47}^{-1} r_{13}^2 T_8^{3/2} \text{ s} . \quad (3)$$

The ionization parameter is $\xi = \beta L_m/r^2 n_e = 10^4 \beta \alpha^{-1} T_8$. In this expression $\beta < 1$ is the ratio of ionizing to MHD luminosity: it is actually the product of two factors, namely the fraction of the MHD wind energy that is randomised in the stand-off shocks, and the fraction of that energy which goes into electrons that radiate in the X-ray band. For a large fraction of the Fe to be hydrogenic, ξ must exceed 10^3 , and this condition would indeed be satisfied unless β were very small. The effective depth d_i to which Fe and other metals can be ionized is given by balancing the number rate of ionizing photons per square centimetre, $\beta L_m/4\pi r^2$ divided by their mean energy (about 10 keV), against the recombination rate $n_{Fe} \delta d_i/t_{rec}$. In this expression, $\delta \gtrsim 1$ takes into account the recombination of other metals besides Fe.

Standard calculations of photoionization of optically-thin slabs (e.g. Young, 1999) show that the equivalent width of the Fe K-alpha line, for solar abundances, is about 0.5 keV; the line is twice as strong, i.e. 1 keV equivalent width, if the Fe has ten times solar abundances (and even stronger for still higher enrichment). These results are applicable in the present context provided only that one further condition is satisfied: namely, that the ionizing photons encounter a Fe ion before being scattered by free electrons – provided, in other words, that $\tau_T = \sigma_T d_i n_e$ does not greatly exceed unity. We find that

$$d_i \sim 3 c t_{rec} x_{Fe}^{-1} (\beta/\alpha \delta) \simeq 6 \times 10^9 (\beta/\alpha^2 \delta) T_8^{3/2} L_{47}^{-1} r_{13}^2 \zeta_{Fe\odot}^{-1} \text{ cm} \quad (4)$$

where $x_{Fe} = 4 \times 10^{-5} \zeta_{Fe\odot}$ is the Fe abundance normalized to the solar value. The Thomson depth through this layer $\tau_T = \sigma_T d_i n_e \simeq 4 \times 10^2 (\beta/\alpha \delta \zeta_{Fe\odot}) T_8^{1/2} \lesssim 3$ provided $(\beta/\alpha \delta \zeta_{Fe\odot}) \lesssim 10^{-2}$. The three parameters β, α and δ in this expression are all somewhat uncertain. However, the relevant material in the funnel and cavity walls comprises the innermost non-collapsed layers of the precursor star so it could well be greatly enriched in Fe (i.e. we might expect $\zeta_{Fe\odot}$ to greatly exceed unity).

Under the foregoing conditions the Fe K- α photon flux is about 0.1 of the X-ray continuum, and is

$$\dot{N}_{LFe} \sim 10^{54} L_{47} \beta \text{ ph/s} \quad (5)$$

The signal observed $t \sim 1.5$ day after the GRB 991216 burst by Piro *et al.* (2000) corresponds, for an assumed distance $D \sim 4$ Gpc, to 6×10^{52} ph/s, and that observed 0.5-1 day after GRB 000214 by Antonelli et al (2000) at $z \sim 0.47$ corresponds to 3×10^{51} ph/s. As is clear from the above discussion, a wind luminosity of 10^{47} erg/s would be sufficient to yield the observed line signal provided that β , the fraction of the power that goes into a photoionizing X-ray continuum, were not below 0.06. β is of course uncertain, but this number does not seem unreasonably high for the efficiency of particle acceleration by relativistic shocks, and suggests that our fiducial value of 10^{47} erg s $^{-1}$ for the overall luminosity need not be an overestimate. The continuum flux from GRB 991216 in the 1-10 keV band is observed to be 50-100 times stronger than the flux in the Fe line at 1.5 days. Since this factor would only be of order 10 for a photoionized slab with the properties envisaged here, this suggests that most of the continuum, even in the X-ray band, could still come from a standard afterglow model, with luminosity declining as a power-law in t , involving a decelerating blast wave. On the other hand, we cannot rule out the possibility that much of the afterglow in all wavebands is due to a continuing power output from a compact remnant. In the latter case, the time-dependence could be more complicated (with possible rapid variability) and such effects should certainly be looked for.

3. Discussion

The total amount of Fe needed to explain the observed K- α line flux, arising in a thin layer of the funnel walls of a collapsar model, amounts to a very modest mass of $M_{Fe} \sim 10^{-8} M_{\odot}$, which could be Fe synthesized in the core. The Fe-enriched core material can easily reach a distance comparable to $r \sim 10^{13}$ cm in 1 day for an expansion velocity below the limit $v \sim 10^9$ cm s $^{-1}$ inferred by Piro *et al.* (2000) (c.f. also Antonelli et al 2000) from the line widths. Such subrelativistic velocities of the envelope material would arise naturally from low Γ shocks propagating through the star following the collapse, as well as from the burst explosion itself at larger angles from the burst jet axis. The natural progenitor scenario in which this can occur is a collapsar or hypernova model (Woosley, 1993, Paczynski, 1998) with “failed supernova” characteristics (i.e. little or no supernova display).

The initial, energetic portion of the relativistic jet, with a typical burst duration of 1 – 10 s, will rapidly expand beyond the stellar envelope, leading in the usual way to shocks and a decelerating blast wave. A continually decreasing fraction of energy, such as put out by a decaying magnetar, may continue being emitted for periods of a day or longer, and its reprocessing by the stellar envelope can be responsible for the observed Fe line emission in GRB 991216. Since the energy in this tail can decay faster than t^{-1} , the usual standard shock gamma-ray and afterglow scenario need not be affected, being determined by the first 1-10 s worth of the energy input.

Finally, a few comments on how the present suggestion contrasts with other possible interpretation of X-ray lines in the afterglow. Because we invoke a continuing power output after the burst (in an MHD wind) the X-ray lines that are observed a day after the burst do not need to

come from material a light day (or more) in extent, as has been suggested earlier (Weth et al 2000, Böttcher, 2000). Such a large radius entails lower densities (and hence slower recombination rates), and thus requires a much larger total mass of Fe. But the most serious problem is that it may require a two-stage event: a supernova explosion, followed by a burst delayed by many days, so as to allow time for a sub-relativistic Fe-rich shell to have reached the requisite distance. Although such a model has been proposed (Stella & Vietri, 1998; Vietri, *et al.* 2000), detailed calculations suggest (cf Böttcher & Fryer 2000) that there are difficulties in understanding how an efficient burst could be generated after such a long time-delay. Such models would have other potential observational consequences which may be used as a test, such as a reddening and flattening of the light curve at late times as inferred in some cases (e.g. Bloom *et al.*, 1999, Galama *et al.*, 1999).

An alternative to the pre-ejection scenario has been proposed by Böttcher & Fryer (2000). They suggest that a very extended torus could be created during the inward-spiralling of a compact object through the envelope of a giant or supergiant – a process that could be a precursor of an exotic type of supernova. This supernova would not only generate the burst, but expel sub-relativistic material which is shock heated when it encounters the torus. The resultant thermal X-ray emission could display line features. Böttcher & Fryer's suggestion is more attractive than the pre-explosion scenario, in that (as in the model proposed here) the X-ray lines come from a region much less than a light day across. However, it may not yield luminosities high enough to explain GRB 991216. More seriously, the shocked supernova ejecta which are postulated to emit the X-rays have a very large optical depth, which reduces the equivalent width of lines and would cause the ionization edge at 9.28 keV to appear as an absorption rather than an emission feature. The large scattering optical depth may even (if it exceeds c/v) inhibit the escape of continuum radiation on a dynamical timescale. However, there could be particular geometries in which a modified scenario along these general lines could meet the constraints.

Further data on X-ray spectral features from afterglows will surely offer important clues to the nature of the precursor star, and the compact object that triggers the burst.

This research has been supported by NASA NAG5-9192, the Guggenheim Foundation, the Sackler Foundation, NSF PHY94-07194 and the Royal Society. We are grateful to A.C. Fabian, L. Piro and a referee for useful comments.

REFERENCES

- Antonelli, L.A., 2000, ApJL in press (astro-ph/0010221)
- Bloom, J.S. *et al.*, 1999, Nature, 401, 453
- Böttcher, M & Fryer, C.L, 2000, ApJ, subm. (astro-ph/0006076)
- Böttcher, M, 2000, ApJ in press (astro-ph/9912030)
- Eichler, D & Levinson, A, 1999, ApJ, 521, L117

- Galama, T, *et al.*, 2000, ApJ 536, 185
- MacFadyen, A & Woosley, S.E., 1999, ApJ 524, 262
- Mészáros , P. & Rees, M.J., 1998, M.N.R.A.S., 299, L10
- Paczynski, B., 1998, ApJ, 494, L45
- Piro, L, *et al.*, 2000, to appear in Nov. 3 issue of Science
- Rees, M.J & Mészáros , P, 1998, ApJ, 496, L1
- Sari, R. & Mészáros , P, 2000, ApJ
- Thompson, C., 1994, MNRAS, 270, 480
- Usov, V.V., 1994, MNRAS, 267, 1035
- Vietri, M & Stella, L.A., 1998, ApJ 507, L45
- Vietri, M, Perola, G, Piro, L & Stella, L, 2000, MNRAS 308, L29.
- Wheeler, J.C, Yi, I, Hoefflich, P & Wang, L, 2000, ApJ in press (astro-ph/9909293)
- Weth, C, Mészáros , P, Kallman, T & Rees, M.J, 2000, ApJ 534, 581
- Woosley, S., 1993, Ap.J., 405, 273
- Young, A.J., 1999, Ph.D. thesis, Cambridge University